

Creating Synergy with **Conventional and Smal Diameter Implants**

NOVEMBER 14, 2017





Brian J. Jackson, DDS

Inside



he demand for implant-supported dentures is projected to increase in coming years, and it is important for clinicians to be aware of the variety of options for patients. In many cases, a patient's bone quality, quantity, and biomechanics, as well as financial means, may call for a blended approach that utilizes conventional-diam-

eter implants (CDIs) and small-diameter implants (SDIs) or mini dental implants (MDIs). When the proper attention is paid to site preservation and careful implant place ment, clinicians can achieve both functional and esthetic success for patients Oral reconstruction supported by osseoin-

tegrated endosseous implants has improved the quality of life for thousands of patients. The demand for implant therapy in the maxillary and mandibular arch is projected to increase as the number of edentulous or partially edentulous patients rises in the fu ture.¹ Research has demonstrated long-term success of an implant-retained overdenture prosthesis within the range of 85% to 99% depending on the specific arch involved.² Predictable outcomes have been demonstrated when bone quality, quantity, and bio-mechanics have served as the foundation of treatment planning.^{3,4}



BRIAN J. JACKSON, DDS Diplomate American Board of Oral Implantol-ogy/Implant Dentistry *Utica, New York*

48 INSIDE DENTISTRY | January 2016 | www.insidedentistry.n

Frequently, clinicians must develop treatment plans for teeth with a poor prognosis, and extraction with site preservation through socket grafting has become a routine procedure.⁵ Site preservation allows for the development of adequate bone, enabling conventional implant placement while es tablishing bone density for initial fixation for SDIs. CDIs are considered when adequate bone

and significant occlusal loads exist or bone grafting procedures demonstrate a favorable long-term prognosis. However, SDIs are an alternative modality when patients present with diminished bone quality, quantity, and reduced biomechanical loads.⁷ Utilizing a va riety of implant designs and diameters based on the patient's existing anatomy, medical conditions and monetary constraints can provide a multitude of treatment options. In this case presentation, a thought-provoking treatment plan focusing on bone qual ity, quantity, age, and biomechanical loads served as the primary considerations in re-solving the patient's chief complaint.



(1.) Panoramic radiograph demonstrated severe bone loss

mandible was anesthetized and teeth Nos.

23, 25, 26, 28, and 30 were removed. The

sockets were debrided with a double-ended

curette and grafted with a mineralized irra-

diated cancellous allograft (Puros", Zimmer

Dental, www.zimmerdental.com) and con-tained with a d-polytetrafluoroethylene (Cytoplast", Osteogenics Biomedical, www. osteogenics.com) barrier. The extractions of

the maxillary arch consisted of teeth Nos. 1, 4,

6, 12, and 13, and the procedure and grafting materials employed in the maxilla mimicked

the mandibular arch (Figure 4). Maxillary

and mandibular removable partial dentures

Case Presentation

A 41-year-old man presented to the office stating that he did not want to wear partials nymore because they were not comfortable The patient's medical history exhibited no significant findings except that he smoked one pack of cigarettes per day. A radiographic survey demonstrated severe horizontal bone loss associated with his maxillary and mandibular teeth (Figure 1). A diagnosis of severe periodontitis was established. The intrao dental examination revealed a total of 16 remaining teeth in the maxillary and man dibular arches (Figure 2). Class II and III periodontal mobility was exhibited by 9 teeth. The patient was wearing a maxillary transiFeedback

tional removable partial denture (Figure 3). At consultation, various treatment plans were presented, including saving specific teeth or full-mouth extraction. In addition, the treatment plans included CDIs and SDIs to enhance support, stability, and retention in prosthetic reconstruction. The agreed upon treatment plan was maxillary and mandibular overdentures. A screw-retained

Inside Implants

superstructure supported by four CDIs would retain the maxillary overdenture. The man-dibular overdenture prosthesis would be retained by four SDIs. The treatment would be performed in a staged approach with extrac-tion and grafting followed by implant placement. Consent, time frame for treatment com-pletion, and provisionalization were reviewed. The pre-implant surgical stage consisted of extraction with site preservation. The

he patient had 16 teeth remaining between the two arches. (3.) The patient's existing lary partial denture. (4.) Frontal view following extractions and grafting. (5.) Maxillary nandibular removable partial dentures were placed. (6.) The surgical template in place mini dental implants in their final position. (8.) Additional extractions were performer nths after implant placement. (9.) The maxillary arch prior to impression taking. (2.) T

(RPDs) were placed, occlusion adjusted, and polished (Figure 5). Surgical Stage

The maxillary implant surgical procedure was initiated 3 months after the extractions and socket grafting. The patient was prepped, draped, and asked to rinse with a 0.12% chlorhexidine mouth rinse for 30 seconds. The blood separation process to develop plate-let-rich plasma was initiated. The patient was anesthetized, and a midcrestal incision with a 15c blade was made and a full mucoperios-teal flap reflected with a periosteal elevator. A surgical guide was placed and four osteotomy "dimples" were made with a surgical long shank No. 4 round bur. The implant surgical protocol drill sequence was 1.3, 2.0, 2.5, and 3.1 mm to a depth of 12 mm for the future 3.8-mm diameter implants. The 4.8-mm diameter implant required two additional drills, 3.8 and 4.1 mm, to complete the osteotomy. Three 3.8-mm x 12-mm RBC Tapered Laser-Lok (BioHorizons, www.biohorizons.com) implants were placed at the crest using a fixture mount and a 2.5mm hex driver. The cover screw was secured to the fixture via a 1.25-mm hex tool. The mu coperiosteal flap was closed using 4.0 vicryl sutures in a horizontal mattress manner.

The mandibular implant surgery was initiated 3 months after the maxillary implants were placed. The preparation and anesthesia of the patient were similar to the maxillary implant surgery. A full mucoperiosteal flap was established with a 15c blade penetrating midcrestally and reflected with a periosteal elevator. A surgical template was placed to aid in the mesial-distal location of the future implant sites (Figure 6). The partial osteotomies were prepared with a 1.1-mm drill to a depth of 6 mm. The four 2-mm x 13-mm O-ball collared MDIs (MDI Mini Dental Implants, 3M Oral Care, www.3mespe.com/implants) were auto-advanced with a finger driver, thumb wrench, and ratchet to final position (Figure 7). The transitional RPD was relieved where the transgingival aspect of the o-ball was located and the occlusion was adjusted. Second stage surgery for the maxillary arch

consisted of the exposure of the conventional implants and extraction of all remaining teeth. This procedure was initiated 6 months post implant placement. Three 3.8-mm x 5-mm and one 4.8-mm x 5-mm titanium healing collars were placed during the surgical un-covering procedure. In addition, teeth Nos. 3, 11, 14, 19, 22, 27, and 30 were extracted (Figure

Inside Implants

8). The transitional RPDs were modified to incorporate additional prosthetic teeth and relined with a soft temporary material. The thetic reconstruction.

Prosthetic Reconstruction The prosthetic reconstruction stage was unique to the specific type of implant design utilized in the treatment plan. The primary objective of the impression stage was to capture the position and angulation of the implants. The maxillary arch employed an open-tray impression technique to capture an abutment level impression (Imprint[®] 3 VPS Impression Material, 3M Oral Care) (Figure 9). The man-dibular MDIs were impressed after placing red" transfers onto the o-ball aspect of the im-plant (Figure 10). A polyvinylsiloxane impression material (Imprint 3) was utilized within a border molded custom tray for both arches. The intraoral impressions were sent to a commercial laboratory where a final work-ing model was established. The maxillary superstructure with locator attachments was designed and manufactured with CAD/

CAM technology (Figure 11). The mandibular


unibase with o-ring housings was made using standard dental laboratory procedures (Figure 12). The passivity of the maxillary superstructure was confirmed intraorally using the Sheffield test and a panoramic radiograph. A maxillary/mandibular relationship utilizing baseplates and wax rims, as well as a mold and en (Figure 13) final prosthesis was used to confirm vertical dimension of occlusion, phonetics, esthetics, and patient acceptance. The final superstructure was fixated to the implants with retaining screws, torqued to 30 Ncm and locator inserts placed with a cumulative retention value of 7.5 lbs (2 pink and 1 blue). The final maxillary and mandibular overdentures were placed 1 week thereafter (Figure 14 and Figure 15).

Discussion Creating synergy with CDIs and SDIs can pro-vide a multitude of treatment alternatives for oral rehabilitation. The size of the implants used depends on various factors of the quality and quantity of bone, as well as biomechanical tress. Site preservation has become a predict able method to optimize bone dimensions and density for future implant sites.85

Conventional implants (3.4 to 6.0 mm) are the primary type of endosseous implants used for long-term predictability in implantretained or supported therapy.10 They can be

utilized when the facial-buccal or facial-lingual dimension exhibits a minimal bone of 6 to 8 mm. Conventional size implants remain the primary option for patients who have adequate bone or who accept bone grafting procedures. and who also have financial means. SDIs (1.8 to 2.9 mm) are an alternative to

Die whon specific criteria are met " The jor reason for utilizing SDIs is minimal width of bone or space between adjacent teeth in a mesio-distal dimension. In addition, SDIs are indicated for patients who are not candidates for invasive bone grafting procedures due to medical reasons, limited finances, or time. SDIs demonstrate high success in the mandibular arch when retaining removable overdentures.12 The maxilla has demonstrated lower success rates than the mandible in regards to implant therapy. Protocols have been established to enhance success rates as they relate to the mandible.¹³ These protocols have centered around minimizing stress to the crestal bone where bone loss is often seen. A critical con-sideration is utilization of implants with a diameter size greater than 3 mm, which reduces stress at the crest of the alveolar bone due to increased surface area. In addition, increasing the number of implants from four to six demonstrates higher success rates via an increase in cumulative surface area. Implant lengths of 12 mm or greater have exhibited two to three





(10.) Red transfers were placed onto the o-ball aspect of the mini dental implants before impressions were made. (11.) The completed maxillary superstructure with locator attachments. (12.) A mandibular unibase was made with o-ring housings to snap over the small-diameter implants. (13.) A maxillary/mandibular relationship was taken utilizing baseplates and wax rims.

52 INSIDE DENTISTRY | January 2016 | www.insidedentistry.net

Inside Implants

times greater success versus 10-mm lengths. Superstructure design without a cantilever component has demonstrated higher overdenture retention with less resultant crestal bone loss. The elimination of cantilevers in the maxillary superstructure bar reduces stress on the distal terminal implants.

Research has demonstrated positive long-term outcomes with the utilization of SDIs in the mandible.14 Rigid fixation of the implant body at the time of placement is crucial for the osseointegrative process to ensue. Secondarily, the auto-advanced technique coupled with a partial osteotomy enhances bicortical stabilization.16 Flapless surgery and immediate load are considerations when utilizing a one-piece SDI.17 However, a full mucoperiosteal flap with a 3-month osseointegrative time period was utilized in this case

In the case report, the decision to utilize conventional implants in the maxilla was based on diminished bone quality and significant biomechanical load. Research has exhibited guarded success rates with maxillary overdentures.¹⁸ SDIs were employed in the mandible because excellent success rates have been demonstrated in similar cases.19 The patient made a treatment decision based on personal finances, as well as an understanding of bone density and biomechanics and how they relate to long-term outcomes.

Conclusion

Oral implantology has become a major discipline in the field of dentistry. Clinicians must evaluate the patient's existing condition and develop thought-provoking treatment plans based on a variety of factors. The utilization

of CDIs and SDIs should be considered in regards to bone, biomechanics, age, and finances. A synergy of different implant modalities can provide alternative approaches to resolve pa tient concerns in a predictable, less invasive, and more economical way.

Disclosure Brian J. Jackson, DDS, has previously lectured for 3M, but did not receive compensation for this article

Acknowledgments The author wishes to acknowledge Tatyana Lyubezhanina, DA, and Caitlin Carparelli, DA, for their assistance in the preparation of this paper, and Matt Weigand, CDT, of Utica Dental Laboratory in Utica, New York, for his dental technical expertise.

References 1. Douglass CW, Shih A, Ostry L. Will there be a need for complete dentures in the United States in 2020? J Prosthet Dent. 2002;87(1):5-8.

2. Goodacre CJ, Bernal G, Rungcharassaeng K, Kan JY. Clinical complications with implants and implant prostheses. J Prosthet Dent. 2003;9(2):121-132. 3. Jemt T, Chai J, Harnett J, et al. A 5-year prospec

tive multicenter follow-up report on overdentures supported by osseointegrated implants. *Int J Oral* Maxillofac Implants. 1996;11(3):291-298. 4. Bergendal T, Enquist B. Implant-supported overdentures: a longitudinal prospective study. Int J Oral

Maxillofac Implants. 1998:13(2):253-262. 5. Sclar AG. Preserving alveolar ridge anatomy following tooth removal in conjunction with immediate implant placement. The Bio-Col technique. Atlas Oral Maxillofac Surg Clin North Am. 1999;7(2):39-59. 6. Nevins R, Mellonig JT. Enhancement of the dam-

aged edentulous ridge to receive dental implants: a combination of allograph and GORE-TEX membrane Int J Periodontics Restorative Dent. 1992;12(2):207-217. 7. Sendax V. Mini Dental Implants: Principles and Practices. St Louis, MO: Mosby; 2013.

8. Jackson BJ, Morcos I. Socket grafting: a predictable technique for site preservation. J Oral Implantol. 2007;33(6):353-364.

9. Irinakis T, Tabesh M. Preserving the socket dimen sions with bone grafting in single sites: an esthetic surgical approach when planning delayed implant placement. J Oral Implantol. 2007;33(3):156-163. 10. Albrektsson T, Zarb G, Worthington P, Eriksson AR. The long-term efficacy of currently used dental implants: a review and proposed criteria of success. *Int* J Oral Maxillofac Implants. 1986;1(1):11-25. 11. English CE, Bonie GC. Diagnostic procedural and

clinical issues with the Sendax mini dental implants. Compend Cont Educ Dent. 2003;24(11 suppl 1):3-25. 12. Shatkin TE, Shatkin S, Oppenheimer BD, Oppenheimer AJ. Mini dental implants for longterm fixed and removable prosthetics: a retrospectiv analysis of 2514 implants placed over a five-year period. Compend Contin Educ Dent. 2007;28(2):92-99.

13. Hutton JE, Heath MR, Chai JY, et al. Factors related to success and failure rates at 3-year follow-up in a multicenter study of overdentures supported by Branemark implants. Int J Oral Maxillofac Implants. 1995;10(1):33-42. 14. Christensen GJ. The 'mini'-implant has arrived. J Am Dent Assoc. 2006;137(3):387-390. 15. Balkin BE, Steflik DE, Naval F. Mini-dental implant

insertion with the auto-advance technique for ongoing applications. J Oral Implantol. 2001;27(1):32-37. 16. Bulard RA, Vance JB. Multi-clinic evaluation using mini-dental implants for long-term denture stabiliza tion: a preliminary biometric evaluation. Compend

Contin Educ Dent. 2005:26(12):892-897. 17. Flanagan D. Flapless dental implant placement. J Oral Implantol. 2007;33(2):75-83. 18. Smedberg JI, Lothigius E, Bodin I, et al. A clinical

and radiological two-year follow-up study of maxillary overdentures on osseointegrated implants. Clin Oral Implants Res. 1993;4(1):39-46. 19. Griffitts TM, Collins CP, Collins PC. Mini dental

implants: an adjunct for retention, stability, and com fort for the edentulous patient. Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2005;100(5):e81-e84



54 INSIDE DENTISTRY | January 2016 | www.insidedentistry.net



To discuss this case, please visit the Customer Portal.

OTHER RECENT CASES

View All »

Prosthetic Design and Fabrication of a Straumann...

July 2, 2018 — Use Cases

As previously described, non-restorable teeth

were removed and grafted. Following integration of the graft, 4 Straumann BLT implants were strategically placed using the Engel protocol. The implants were allowed to...

🦄 by Timothy Kosinski

READ MORE

Cone Beam Computed Tomography in Implant Dentistry:...

May 17, 2018 — Use Cases

In implant dentistry, three-dimensional (3D) imaging can be realised by dental cone beam computed tomography (CBCT), offering volumetric data on jaw bones and teeth with relatively low radiation doses and...

by GreenCT

READ MORE

Periapical Radiograph Not Diagnostic – Power of...

May 15, 2018 — Use Cases

Patient came in with pain, upper left posterior molar... a periapical was inconclusive... no swelling, no fistula. How can we determine the proper treatment plan today without CBCT ..? An essential...



by Scott Ganz

READ MORE

Not a Member yet? Sign up for Free

GET STARTED



The primary mission of Green CT[®] is to aid dental professionals and provide excellent care in a safe environment that promotes healing.

ABOUT US

DENTISTRY UPDATES Green CT Newsletter The Green CT newsletter provides clinicians with the latest news and information from Green CT and Vatech. Sign up below with your email address today! **EMAIL ADDRESS** SIGN UP in

OUR MISSION

WHY GREENCT? CASE STUDIES TRAINING & LEARNING VIDEOS

PRODUCTS

GREEN CT 2

EZ3D-i

RADIATION COMPARISON SENSOR TECHNOLOGY

vatech

2200 Fletcher Ave, Suite 705A Fort Lee, NJ 07024

T. 201.210.5118 F. 201.601.0859 marketing@vatechamerica.com

SUPPORT

ABOUT GREENCT

ABOUT US **NEWS & EVENTS** CONTACT US

VATECH ASSURANCE FAQ CONTACT SUPPORT

REGISTER

ВАСК ТО ТОР ∧

Terms Of Use|Privacy Statement